ELSEVIER

www.elsevier.nl/locate/jorganchem

Journal of Organometallic Chemistry 595 (2000) 315–316

Journal of Organometallic Chemistry

Author Index of Volume 595

Abd El Baky, C., 285
Adams, H., 199
Adamus, G., 66
Ahlgrén, M., 232
Albert, J., 54

Balavoine, G.G.A., 261
Bandini, A.L., 224
Banditelli, G., 224
Banerjee, S., 140
Bates, T.F., 87
Bayler, A., 296
Beckett, M.A., 241
Berglund Åhman, A., 102
Blaurock, S., 126
Böhm, V.P.W., 186
Bosque, R., 54
Buika, G., 66

Canty, A.J., 296
Castro, A., 36
Che, C.-M., 114
Cheung, K.-K., 114
Choi, M.-Y., 114
Cotton, F.A., 300

Dahrouch, M., 153
Dallemer, F., 261
Dandekar, S.A., 87
Daran, J.-C., 261
Deeney, T., 191

Eisen, M.S., 126
Eriksson, H., 102

Fawzi, R., 166
Fernández, A., 199
Fernández, J.J., 199
Fey, O., 285
Fredin, L., 248

Galakhov, M.V., 36
Galan-Fereres, M., 126
García-Vázquez, J.A., 59
Gelbrich, T., 21
Gelinsky, M., 140
Ghose, S., 140
Gómez, M., 36
Gómez-Sal, P., 36
Gornitzka, H., 153
Gouygou, M., 261
Granell, J., 54
Gray, G., 191
Gražulevičius, J.V., 66
Grobelny, Z., 66
Grün, K., 285
Gstöttmayr, C.W.K., 186
Guerchais, V., 81

Håkansson, M., 102
Hecht, E., 21
Henkel, G., 166
Herrmann, W.A., 186
Hey-Hawkins, E., 126
Hibbs, D.E., 241
Hobbold, M., 12
Hogarth, G., 134
Horner, G.M., 215
Hor, T.S.A., 276
Hourihane, R., 191
Huang, X., 268
Hursthouse, M.B., 241
Hu, Z., 268
Hydrio, J., 261

Jagner, S., 102
Jankowski, S., 109
Jeges, G., 93
Jeske, J., 12
Jiang, C., 276
Jones, P.G., 12
Jung, J., 147

Keglevich, G., 109
Kégl, T., 93
Keun Park, S., 147
Khabashesku, V.N., 248
Kiesewetter, J., 81
Kim, H., 147
Koch, T., 126
Kollár, L., 93
Koshino, H., 208
Kudin, K.N., 248
Kumar, G.R., 140
Kyun Noh, S., 147

Laitinen, R.S., 232
Lam, H.-Y., 70
Lavender, M.H., 134
Lee, D.-h., 147
Lee, F.-W., 114
Lee, S.-M., 70
Lindner, E., 166
Liu, H., 276
Liu, L.K., 276
Longlet, J.J., 87
López-Torres, M., 199

Malik, K.M.A., 241
Malisch, W., 285
Margrave, J.L., 248
Martín, A., 36
Mathur, P., 140
Maynard, N.P., 215
Mereiter, K., 93
Minghetti, G., 224
Miyaura, N., 31
Mohr, M., 166
Mok, K.F., 276
Moszner, M., 178

Nachtigal, C., 166
Nakata, T., 208
Nieuwenhuyzen, M., 292

Obora, Y., 1
Oilunkaniemi, R., 232
Ortigueira, J.M., 199
Owen, P., 241

Pan, Y., 268
Pérez-Lourido, P., 59
Petőlz, G., 93
Philip, R., 140
Plat, D., 126
Poignant, G., 81

Rickard, C.E.F., 215
Rivière-Baudet, M., 153
Rombach, M., 140
Romero, J., 59
Rudzinski, J., 109
Russo, U., 59

Sánchez, F., 36
Satake, A., 208
Saunders, G.C., 292
Shukri, K., 134
Sieler, J., 21
Skelton, B.W., 296
Sousa, A., 59
Spalding, T., 191
Stiriba, S.-E., 300
Stolarzewicz, A., 66
Streubel, R., 12
Sturm, T., 93
Suárez, A., 199
Sun, H., 268
Szelke, H., 109

PII: S0022-328X(99)00718-4

Tanaka, M., 1
Tavera, R., 54
Teng, X., 268
Thiele, K.-H., 21
Thomas, R.D., 87
Trivedi, R., 140
Trösch, A., 178

Ueda, M., 31

Vahrenkamp, H., 140, 178
Varma, K.S., 241
Vázquez-García, D., 199
Vila, J.M., 199

Wang, Z.-X., 158
Weissensteiner, W., 93
Weskamp, T., 186
White, A.H., 296
Wołowiec, S., 178
Wong, W.-Y., 70
Woodgate, P.D., 215
Wood, K.A., 87

Yan, Y.K., 276
Yeo, J.S.L., 276
Yokochi, A., 300

Zhao, H., 158
Zhou, X.-G., 158
Zhou, Z.-Y., 158
Zubieta, J., 59

ELSEVIER

www.elsevier.nl/locate/jorganchem

Journal of Organometallic Chemistry 595 (2000) 317–322

Journal of Organometallic Chemistry

Subject Index of Volume 595

Alkenyl ligand
Reactions of $[\{\mu\text{-}\eta^2\text{-}C(R^1)\text{=}CHR^2\}(\mu\text{-}CO)Fe_2(CO)_6]^-$ with $(\mu\text{-}S_2)Fe_2(CO)_6$: synthesis and structures of $[\mu\text{-}\eta^2\text{-}C(R^1)\text{=}CHR^2](\mu\text{-}RS)[Fe_2(CO)_6]_2(\mu_4\text{-}S)$ (R = alkyl, phenyl or acyl), 158

Alkoxides
Monomeric organoaluminium complexes $RAl(OR^*)_2$ and R_2AlOR^* with an optically active amino alkoxide ligand, 21

Alkoxysilanes
Regiospecific synthesis of alpha-lithiated alkoxysilanes, 87

Alkylidene
Binuclear hydrido platinum(II) complexes: syntheses of $[\{Pt(P\text{–}P)\}_2(\mu\text{-}CHCH_2R)(\mu\text{-}H)][BF_4]$ (R = C_6H_5, H) and easy cleavage of a P–C bond in a chelating 1,4-bis (diphenyl)phosphinobutane (P–P), 224

Alkyne
Synthesis, electrochemistry and structural characterization of new dimeric cobalt η^2-alkyne carbonyl complexes, 70

Alkynyl complexes
Selectivity in alkynylation: the reaction between $Ru_2(LL)_4Cl$ and $Me_3SnC\text{≡}CR$ (LL = 2-anilinopyridine, 2-chloro- and 2-bromo-oxypyridine), 300

Alpha-lithiated alkoxysilanes
Regiospecific synthesis of alpha-lithiated alkoxysilanes, 87

Alpha-lithiation
Regiospecific synthesis of alpha-lithiated alkoxysilanes, 87

Alpha-silyl carbanions
Regiospecific synthesis of alpha-lithiated alkoxysilanes, 87

Aluminium
Monomeric organoaluminium complexes $RAl(OR^*)_2$ and R_2AlOR^* with an optically active amino alkoxide ligand, 21

Amines
Monomeric organoaluminium complexes $RAl(OR^*)_2$ and R_2AlOR^* with an optically active amino alkoxide ligand, 21

Aminocarbene complexes
Synthese und Struktur von {[*N*-[Bis(diisopropylamino)phosphanyl]amino(phenyl)carben}dicarbonyl(η^5 - methylcyclopentadienyl)mangan(0)} und versuchte Darstellung von {[2-Bis(trimethylsilyl)methyl-3-phenyl-2*H*-azaphosphiren-κ*P*]dicarbonyl(η^5-methylcyclopentadienyl)mangan(0)}, 12

Aminophosphines
Preparation, properties and reactions of metal-containing heterocycles: Part C: tetraazatetraphosphadimolybdacyclophanes: synthesis, isolation, characterization, and X-ray crystal structures, 166

Aromatic bridge
Convenient routes to mono- and dinuclear (C_5Me_5)iron(II) complexes bearing acyl, alkynyl, aryl and thienyl ligands, 81

Aromatic ditellurides
The X-ray crystallographic study of the reaction of bis(2-thienyl)ditelluride with tetrakis(triphenylphosphine)platinum or -palladium, 232

Aryloxy–zirconocene complex
Dehydrocoupling polymerization of arylsilanes with chloro(aryloxy)bis(cyclopentadienyl)zirconium complex catalysts, 1

Arylsilanes
Dehydrocoupling polymerization of arylsilanes with chloro(aryloxy)bis(cyclopentadienyl)zirconium complex catalysts, 1

Binuclear complexes
Binuclear hydrido platinum(II) complexes: syntheses of $[\{Pt(P\text{–}P)\}_2(\mu\text{-}CHCH_2R)(\mu\text{-}H)][BF_4]$ (R = C_6H_5, H) and easy cleavage of a P–C bond in a chelating 1,4-bis (diphenyl)phosphinobutane (P–P), 224

Bis(trimesitylgermyl)-carbodiimide
Bis(trimesitylgermylcarbodiimido)germylene, trimesitylgermylcyanamide and trimesitylgermylcarbodiimide, 153

Bis(trimesitylgermylcarbodiimido)germylene
Bis(trimesitylgermylcarbodiimido)germylene, trimesitylgermylcyanamide and trimesitylgermylcarbodiimide, 153

Borane adduct
P–H functionalised phosphinoalkoxides as ligands: synthesis and catalytic properties of $[Cp_2Zr(Me)(cyclo\text{-}1\text{-}O\text{-}2\text{-}PH(Tipp)\text{-}C_6H_{10})]$ (Tipp = 2,4,6-$Pr^i_3C_6H_2$), $[Cp_2Zr(Me)(cyclo\text{-}1\text{-}O\text{-}2\text{-}PH(Tipp)\{Mo(CO)_5\}\text{-}C_6H_{10})]$ and $[Cp_2Zr(BH_4)(cyclo\text{-}1\text{-}O\text{-}2\text{-}PH(Mes)(BH_3)\text{-}C_6H_{10})]$ (Mes = 2,4,6-$Me_3C_6H_2$), and molecular structure of $(R_C,R_C,R_P/S_C,S_C,S_P)$-$[Cp_2Zr(Me)(cyclo\text{-}1\text{-}O\text{-}2\text{-}PH(Tipp)\text{-}C_6H_{10})]$, 126

Borasiloxanes
cyclo-Boratrisiloxane and *cyclo*-diboratetrasiloxane derivatives and their reactions with amines: crystal and molecular structure of $(p\text{-}BrC_6H_4BO)_3(Ph_2SiO)_2$, 241

Boron
cyclo-Boratrisiloxane and *cyclo*-diboratetrasiloxane derivatives and their reactions with amines: crystal and molecular structure of $(p\text{-}BrC_6H_4BO)_3(Ph_2SiO)_2$, 241

Synthesis of dioxazaborocines from *N,N'*-alkylbridged-bis(bis(2-hydroxybenzyl)aminomethyl)amines, 215

Bridging ligand
Synthesis and spectroscopic characterisation of compounds with formula $[\{Fe_2S_2(CO)_5\}Ph_2P(CH_2)_nPPh_2\{Fe(CO)_4\}]$ and $[\{Fe_2S_2(CO)_5\}_2Ph_2P(CH_2)_nPPh_2]$ (n = 3, 4 and 6), 191

Bridging-sulfur ligand
Reactions of $[\{\mu\text{-}\eta^2\text{-}C(R^1)\text{=}CHR^2\}(\mu\text{-}CO)Fe_2(CO)_6]^-$ with $(\mu\text{-}S_2)Fe_2(CO)_6$: synthesis and structures of $[\mu\text{-}\eta^2\text{-}C(R^1)\text{=}CHR^2](\mu\text{-}RS)[Fe_2(CO)_6]_2(\mu_4\text{-}S)$ (R = alkyl, phenyl or acyl), 158

Carbazole
Electron-transfer processes in the reaction of K^-, $K^+(15\text{-crown-}5)_2$ with carbazole: unexpected formation of metallic potassium, 66

Carbene
Convenient routes to mono- and dinuclear (C_5Me_5)iron(II) complexes bearing acyl, alkynyl, aryl and thienyl ligands, 81

Carbon–carbon bond formation
N-Heterocyclic carbenes: Part 26. *N*-Heterocyclic carbene complexes of palladium(0): synthesis and application in the Suzuki cross-coupling reaction, 186

PII: S0022-328X(99)00719-6

Carbon disulfide
Reactivity of $[Fe_2(CO)_6(\mu\text{-}CO)(\mu\text{-}dppm)]$ towards unsaturated sulfur-containing compounds, SO_2, CS_2 and ArNCS: facile carbon–sulfur bond cleavage at the diiron centre, 134
Carbon–fluorine bond activation
Regioselective C–F and C–H bond activation: synthesis and structure of an asymmetric rhodium complex comprising a η^5:η^1:η^1-cyclopentadienyl-bis(phosphine) ligand, 292
Carbon–hydrogen bond activation
Regioselective C–F and C–H bond activation: synthesis and structure of an asymmetric rhodium complex comprising a η^5:η^1:η^1-cyclopentadienyl-bis(phosphine) ligand, 292
Carbonylation
Platinum complexes of heteroannularly bridged heterobidentate ferrocenyl diphosphine ligands: their molecular structure and their use in catalytic carbonylation reactions, 93
Chirality
Monomeric organoaluminium complexes $RAl(OR^*)_2$ and R_2AlOR^* with an optically active amino alkoxide ligand, 21
Cobalt
Synthesis, electrochemistry and structural characterization of new dimeric cobalt η^2-alkyne carbonyl complexes, 70
Complex
Synthesis, optical nonlinearity and crystal structure of a novel mixed-chalcogen, thiolato-bridged complex $[(\eta^5\text{-}C_5H_5)_2Mo_2(\mu\text{-}S)(\mu\text{-}Te)(\mu\text{-}SPh)_2]$, 140
Complexes
Reactions of $[\{\mu\text{-}\eta^2\text{-}C(R^1)\text{=}CHR^2\}(\mu\text{-}CO)Fe_2(CO)_6]^-$ with $(\mu\text{-}S_2)Fe_2(CO)_6$: synthesis and structures of $[\mu\text{-}\eta^2\text{-}C(R^1)\text{–}CHR^2](\mu\text{-}RS)[Fe_2(CO)_6]_2(\mu_4\text{-}S)$ (R = alkyl, phenyl or acyl), 158
Coupling reaction
Selectivity in alkynylation: the reaction between $Ru_2(LL)_4Cl$ and $Me_3SnC\text{≡}CR$ (LL = 2-anilinopyridine, 2-chloro- and 2-bromo-oxypyridine), 300
Cross coupling
N-Heterocyclic carbenes: Part 26. *N*-Heterocyclic carbene complexes of palladium(0): synthesis and application in the Suzuki cross-coupling reaction, 186
Cryogenic reactions
Matrix isolation infrared and density functional theoretical studies of cryogenic reactions of silicon atoms with dimethyl ether and methanol: new route to generation and stabilization of transient silanones, 248
Crystal structure
Diorganotin(IV) derivatives of arenephosphinothiol ligands. The crystal structure of $[Ph_2Sn\{2\text{-}(Ph_2P)C_6H_4S\}_2]$ and $[Me_2Sn\{2\text{-}(Ph_2PO)\text{-}6\text{-}(Me_3Si)C_6H_3S\}_2]$, 59
Organopalladium(IV) complexes containing phosphine ligands, and the structure of the platinum(IV) complex $[PtMe_3(bpy)(PPh_3)][O_3SCF_3]$ (bpy = 2,2′-bipyridine), 296
Crystal structures
Isocyanide ligation at ruthenium(II) complexes with chelating tertiary amine and porphyrin ligands. Structural and electrochemical studies, 114
Synthesis and structure of Rh(I) tris(pyrazolyl)borate complexes of the type $Tp^{Ph,Me}RhL_1L_2$, 178
Synthesis, electrochemistry and structural characterization of new dimeric cobalt η^2-alkyne carbonyl complexes, 70
[2 + 2]-Cycloaddition
[2 + 2]-Cycloaddukte PH-funktioneller Phosphenium-Komplexe mit Alkylisothiocyanaten: Darstellung von $C_5R_5(OC)_2$<ovl type = "tsqb">M–P(H)(*t*-Bu)–C(=NR′)–S (R = H, Me; M = Mo, W; R′ = Me, Et, *t*-Bu) und Reaktion unter Beanspruchung der PH-Funktion: Phosphenium-Übergangsmetallkomplexe, 37, 285
Cyclometallated
New cyclometallated platinum(II) compounds with thiosemicarbazones: crystal and molecular structure of $[Pt\{4\text{-}MeC_6H_3C(Me)\text{=}NN\text{–}C(S)NH_2\}(PPh_3)]$, 199
Cyclometallation
Cyclometallation of anthracen-9-ylmethylene-phenyl-amine by palladium(II) compounds, 54

Dehydrocoupling polymerization
Dehydrocoupling polymerization of arylsilanes with chloro(aryloxy)bis(cyclopentadienyl)zirconium complex catalysts, 1
Density functional calculations
Matrix isolation infrared and density functional theoretical studies of cryogenic reactions of silicon atoms with dimethyl ether and methanol: new route to generation and stabilization of transient silanones, 248
Dimeric carbonyl
Reactivity of $[Fe_2(CO)_6(\mu\text{-}CO)(\mu\text{-}dppm)]$ towards unsaturated sulfur-containing compounds, SO_2, CS_2 and ArNCS: facile carbon–sulfur bond cleavage at the diiron centre, 134
Dimolybdacyclophanes
Preparation, properties and reactions of metal-containing heterocycles: Part C: tetraazatetraphosphadimolybdacyclophanes: synthesis, isolation, characterization, and X-ray crystal structures, 166
Dinuclear
Convenient routes to mono- and dinuclear (C_5Me_5)iron(II) complexes bearing acyl, alkynyl, aryl and thienyl ligands, 81
Dioxazaborocine
Synthesis of dioxazaborocines from *N*,*N*′-alkylbridged-bis(bis(2-hydroxybenzyl)aminomethyl)amines, 215
Diphosphenes
Synthese und Struktur von {{*N*-[Bis(diisopropylamino)-phosphanyl]amino(phenyl)carben}dicarbonyl(η^5 - methylcyclopentadienyl)mangan(0)} und versuchte Darstellung von {[2-Bis(trimethylsilyl)methyl-3-phenyl-2*H*-azaphosphiren-κ*P*]dicarbonyl(η^5-methylcyclopentadienyl)mangan(0)}, 12
Diphosphine
Synthesis and spectroscopic characterisation of compounds with formula $[\{Fe_2S_2(CO)_5\}Ph_2P(CH_2)_nPPh_2\{Fe(CO)_4\}]$ and $[\{Fe_2S_2(CO)_5\}_2Ph_2P(CH_2)_nPPh_2]$ (n = 3, 4 and 6), 191
Diruthenium
Selectivity in alkynylation: the reaction between $Ru_2(LL)_4Cl$ and $Me_3SnC\text{≡}CR$ (LL = 2-anilinopyridine, 2-chloro- and 2-bromo-oxypyridine), 300
Disiloxane-bridged metallocene
Synthesis and characterization of Group 4 metallocene complexes with two disiloxanediyl bridges, 147
Doubly bridged metallocene
Synthesis and characterization of Group 4 metallocene complexes with two disiloxanediyl bridges, 147

Electrochemistry
Isocyanide ligation at ruthenium(II) complexes with chelating tertiary amine and porphyrin ligands. Structural and electrochemical studies, 114
Electron transfer
Electron-transfer processes in the reaction of K^-, K^+ (15-crown-5)$_2$ with carbazole: unexpected formation of metallic potassium, 66

Ferrocene
Platinum complexes of heteroannularly bridged heterobidentate ferrocenyl diphosphine ligands: their molecular structure and their use in catalytic carbonylation reactions, 93
Ferrocenyl
Synthesis, electrochemistry and structural characterization of new dimeric cobalt η^2-alkyne carbonyl complexes, 70

Fluorene
Synthesis, electrochemistry and structural characterization of new dimeric cobalt η^2-alkyne carbonyl complexes, 70
Fragmentation
Evidence for the involvement of pentacoordinate P-intermediates during the UV light-mediated fragmentation of a 2-phosphabicyclo[2.2.2]octene 2-oxide in the presence of O-nucleophiles, 109

Hafnium
Synthesis and characterization of Group 4 metallocene complexes with two disiloxanediyl bridges, 147
2*H*-azaphosphirene complexes
Synthese und Struktur von {{*N*-[Bis(diisopropylamino)-phosphanyl]amino(phenyl)carben}dicarbonyl(η^5 - methylcyclopentadienyl)mangan(0)} und versuchte Darstellung von {[2-Bis(trimethylsilyl)methyl-3-phenyl-2*H*-azaphosphiren-κ*P*]dicarbonyl(η^5-methylcyclopentadienyl) mangan(0)}, 12
Heterodinuclear molybdenum phosphine complex
P–H functionalised phosphinoalkoxides as ligands: synthesis and catalytic properties of [$Cp_2Zr(Me)$(*cyclo*-1-O-2-PH(Tipp)-C_6H_{10})] (Tipp = 2,4,6-$Pr^i_3C_6H_2$), [$Cp_2Zr(Me)$(*cyclo*-1-O-2-PH(Tipp){$Mo(CO)_5$}–C_6H_{10})] and [$Cp_2Zr(BH_4)$(*cyclo*-1-O-2-PH(Mes)(BH_3)–C_6H_{10})] (Mes = 2,4,6-$Me_3C_6H_2$), and molecular structure of (R_C,R_C,R_P/S_C,S_C,S_P)-[$Cp_2Zr(Me)$-(*cyclo*-1-O-2-PH(Tipp)–C_6H_{10})], 126
Heterometallic
Reactions of [$Pt_2(\mu\text{-}S)_2(PPh_3)_4$] with Group 6 and 7 metal carbonyls; crystal structure of the apparently unsaturated heterometallic complex [$Pt_2Re(\mu_3\text{-}S)_2(PPh_3)_4(CO)_3]^+$, 276
High dilution
Preparation, properties and reactions of metal-containing heterocycles: Part C: tetraazatetraphosphadimolybdacyclophanes: synthesis, isolation, characterization, and X-ray crystal structures, 166
2H-1,3-Naphthoxazine
Synthesis of dioxazaborocines from *N*,*N*′-alkylbridged-bis(bis(2-hydroxybenzyl)aminomethyl)amines, 215
Hydride
Binuclear hydrido platinum(II) complexes: syntheses of [{Pt(P–P)}$_2$(μ-$CHCH_2R$)(μ-H)][BF_4] (R = C_6H_5, H) and easy cleavage of a P–C bond in a chelating 1,4-bis (diphenyl)phosphinobutane (P–P), 224
Hydroformylation
Platinum complexes of heteroannularly bridged heterobidentate ferrocenyl diphosphine ligands: their molecular structure and their use in catalytic carbonylation reactions, 93

Imido η^2-iminoacyl, η^2-acyl cyclopentadienyl complexes
Chemical behaviour of alkyl imido cyclopentadienyl niobium and tantalum(V) complexes in insertion processes. X-ray crystal structures of [MCpCl(NAr){η^2-C(Me)=NAr}] (Ar = 2,6-$Me_2C_6H_3$; M = Nb, Cp = η^5-$C_5H_4SiMe_3$; M = Ta, Cp = η^5-C_5Me_5) and [Ta(η^5-C_5Me_5)Me(NAr){η^2-$C(CH_2CMe_2Ph)$=O}] (Ar = 2,6-$Me_2C_6H_3$), 36
Infrared spectroscopy
Matrix isolation infrared and density functional theoretical studies of cryogenic reactions of silicon atoms with dimethyl ether and methanol: new route to generation and stabilization of transient silanones, 248
Intramolecular proton transfer
Electron-transfer processes in the reaction of K^-, K^+(15-crown-5)$_2$ with carbazole: unexpected formation of metallic potassium, 66

Iron
Convenient routes to mono- and dinuclear (C_5Me_5)iron(II) complexes bearing acyl, alkynyl, aryl and thienyl ligands, 81
Reactions of [{μ-η^2-$C(R^1)$=CHR^2}(μ-CO)$Fe_2(CO)_6$]$^-$ with (μ-S_2)$Fe_2(CO)_6$: synthesis and structures of [μ-η^2-$C(R^1)$=CHR^2](μ-RS)[$Fe_2(CO)_6$]$_2$(μ_4-S) (R = alkyl, phenyl or acyl), 158
Reactivity of [$Fe_2(CO)_6$(μ-CO)(μ-dppm)] towards unsaturated sulfur-containing compounds, SO_2, CS_2 and ArNCS: facile carbon–sulfur bond cleavage at the diiron centre, 134
Synthesis and spectroscopic characterisation of compounds with formula [{$Fe_2S_2(CO)_5$}$Ph_2P(CH_2)_nPPh_2${$Fe(CO)_4$}] and [{$Fe_2S_2(CO)_5$}$_2Ph_2P(CH_2)_nPPh_2$] (n = 3, 4 and 6), 191
The thermal rearrangement of –[Me_2Si-η^5-$C_5H_4(CO)$-$Fe(CO)_2Fe(CO)$-η^5-$C_5H_4SiMe_2$]– in the presence of phosphorus ligands, 268
Isocyanide complexes
Isocyanide ligation at ruthenium(II) complexes with chelating tertiary amine and porphyrin ligands. Structural and electrochemical studies, 114
Isomer
Synthesis and spectroscopic characterisation of compounds with formula [{$Fe_2S_2(CO)_5$}$Ph_2P(CH_2)_nPPh_2${$Fe(CO)_4$}] and [{$Fe_2S_2(CO)_5$}$_2Ph_2P(CH_2)_nPPh_2$] (n = 3, 4 and 6), 191

Lithium
Regiospecific synthesis of alpha-lithiated alkoxysilanes, 87

Macrocyclization
Preparation, properties and reactions of metal-containing heterocycles: Part C: tetraazatetraphosphadimolybdacyclophanes: synthesis, isolation, characterization, and X-ray crystal structures, 166
Manganese
Synthese und Struktur von {{*N*-[Bis(diisopropylamino)-phosphanyl]amino(phenyl)carben}dicarbonyl(η^5 - methylcyclopentadienyl)mangan(0)} und versuchte Darstellung von {[2-Bis(trimethylsilyl)methyl-3-phenyl-2*H*-azaphosphiren-κ*P*]dicarbonyl(η^5-methylcyclopentadienyl) mangan(0)}, 12
Matrix isolation
Matrix isolation infrared and density functional theoretical studies of cryogenic reactions of silicon atoms with dimethyl ether and methanol: new route to generation and stabilization of transient silanones, 248
Mechanism
Evidence for the involvement of pentacoordinate P-intermediates during the UV light-mediated fragmentation of a 2-phosphabicyclo[2.2.2]octene 2-oxide in the presence of O-nucleophiles, 109
Metallocene complexes
Synthesis and characterization of Group 4 metallocene complexes with two disiloxanediyl bridges, 147
Metalloligand
Reactions of [$Pt_2(\mu\text{-}S)_2(PPh_3)_4$] with Group 6 and 7 metal carbonyls; crystal structure of the apparently unsaturated heterometallic complex [$Pt_2Re(\mu_3\text{-}S)_2(PPh_3)_4(CO)_3]^+$, 276
Metathesis
The thermal rearrangement of –[Me_2Si-η^5-$C_5H_4(CO)Fe(CO)_2$-$Fe(CO)$-η^5-$C_5H_4SiMe_2$]– in the presence of phosphorus ligands, 268
Molybdenum
Synthesis, optical nonlinearity and crystal structure of a novel mixed-chalcogen, thiolato-bridged complex [(η^5-C_5H_5)$_2$-$Mo_2(\mu\text{-}S)(\mu\text{-}Te)(\mu\text{-}SPh)_2$], 140

Molybdenum complexes
Preparation, properties and reactions of metal-containing heterocycles: Part C: tetraazatetraphosphadimolybdacyclophanes: synthesis, isolation, characterization, and X-ray crystal structures, 166
Mössbauer
Synthesis and spectroscopic characterisation of compounds with formula $[\{Fe_2S_2(CO)_5\}Ph_2P(CH_2)_nPPh_2\{Fe(CO)_4\}]$ and $[\{Fe_2S_2(CO)_5\}_2Ph_2P(CH_2)_nPPh_2]$ ($n = 3$, 4 and 6), 191

N-arylsulfonyl aldimines
Rhodium-catalyzed phenylation of *N*-arylsulfonyl aldimines with sodium tetraphenylborate or trimethyl(phenyl)stannane, 31
N-Heterocyclic carbenes
N-Heterocyclic carbenes: Part 26. *N*-Heterocyclic carbene complexes of palladium(0): synthesis and application in the Suzuki cross-coupling reaction, 186
Niobium
Chemical behaviour of alkyl imido cyclopentadienyl niobium and tantalum(V) complexes in insertion processes. X-ray crystal structures of $[MCpCl(NAr)\{\eta^2\text{-}C(Me)\text{=}NAr\}]$ ($Ar = 2,6\text{-}Me_2C_6H_3$; $M = Nb$, $Cp = \eta^5\text{-}C_5H_4SiMe_3$; $M = Ta$, $Cp = \eta^5\text{-}C_5Me_5$) and $[Ta(\eta^5\text{-}C_5Me_5)Me(NAr)\{\eta^2\text{-}C(CH_2CMe_2Ph)\text{=}O\}]$ ($Ar = 2,6\text{-}Me_2C_6H_3$), 36
NMR spectroscopy
Synthesis and structure of Rh(I) tris(pyrazolyl)borate complexes of the type $Tp^{Ph,Me}RhL_1L_2$, 178
The X-ray crystallographic study of the reaction of bis(2-thienyl)-ditelluride with tetrakis(triphenylphosphine)platinum or -palladium, 232
^{15}N-NMR
Synthesis and structural analysis of palladium(II) pyridinylpyrazole complexes by ^{1}H-, ^{13}C-, ^{15}N-NMR and X-ray diffraction. Comparison of binuclear methylpalladium, chloromethylpalladium, and dichloropalladium complexes by ^{15}N-NMR, 208
Nuclear magnetic resonance
Platinum complexes of heteroannularly bridged heterobidentate ferrocenyl diphosphine ligands: their molecular structure and their use in catalytic carbonylation reactions, 93
Nucleophility
Evidence for the involvement of pentacoordinate P-intermediates during the UV light-mediated fragmentation of a 2-phosphabicyclo[2.2.2]octene 2-oxide in the presence of O-nucleophiles, 109

Organotin complexes
Diorganotin(IV) derivatives of arenephosphinothiol ligands. The crystal structure of $[Ph_2Sn\{2\text{-}(Ph_2P)C_6H_4S\}_2]$ and $[Me_2Sn\text{-}\{2\text{-}(Ph_2PO)\text{-}6\text{-}(Me_3Si)C_6H_3S\}_2]$, 59
Oxopyridine
Selectivity in alkynylation: the reaction between $Ru_2(LL)_4Cl$ and $Me_3SnC\equiv CR$ (LL = 2-anilinopyridine, 2-chloro- and 2-bromo-oxypyridine), 300

Palladium
Cyclometallation of anthracen-9-ylmethylene-phenyl-amine by palladium(II) compounds, 54
N-Heterocyclic carbenes: Part 26. *N*-Heterocyclic carbene complexes of palladium(0): synthesis and application in the Suzuki cross-coupling reaction, 186
Organopalladium(IV) complexes containing phosphine ligands, and the structure of the platinum(IV) complex $[PtMe_3(bpy)(PPh_3)][O_3SCF_3]$ (bpy = 2,2′-bipyridine), 296
Platinum complexes of heteroannularly bridged heterobidentate ferrocenyl diphosphine ligands: their molecular structure and their use in catalytic carbonylation reactions, 93
Synthesis and structural analysis of palladium(II) pyridinylpyrazole complexes by ^{1}H-, ^{13}C-, ^{15}N-NMR and X-ray diffraction. Comparison of binuclear methylpalladium, chloromethylpalladium, and dichloropalladium complexes by ^{15}N-NMR, 208
Palladium complex
The X-ray crystallographic study of the reaction of bis(2-thienyl)ditelluride with tetrakis(triphenylphosphine)platinum or -palladium, 232
P–C cleavage
Binuclear hydrido platinum(II) complexes: syntheses of $[\{Pt(P\text{–}P)\}_2(\mu\text{-}CHCH_2R)(\mu\text{-}H)][BF_4]$ ($R = C_6H_5$, H) and easy cleavage of a P–C bond in a chelating 1,4-bis (diphenyl)phosphinobutane (P–P), 224
Pentacoordinated intermediate
Evidence for the involvement of pentacoordinate P-intermediates during the UV light-mediated fragmentation of a 2-phosphabicyclo[2.2.2]octene 2-oxide in the presence of O-nucleophiles, 109
P–H functionalised phosphinoalkoxide
P–H functionalised phosphinoalkoxides as ligands: synthesis and catalytic properties of $[Cp_2Zr(Me)(\textit{cyclo}\text{-}1\text{-}O\text{-}2\text{-}PH(Tipp)\text{-}C_6H_{10})]$ ($Tipp = 2,4,6\text{-}Pr^i_3C_6H_2$), $[Cp_2Zr(Me)(\textit{cyclo}\text{-}1\text{-}O\text{-}2\text{-}PH(Tipp)\{Mo(CO)_5\}\text{-}C_6H_{10})]$ and $[Cp_2Zr(BH_4)(\textit{cyclo}\text{-}1\text{-}O\text{-}2\text{-}PH(Mes)(BH_3)\text{-}C_6H_{10})]$ ($Mes = 2,4,6\text{-}Me_3C_6H_2$), and molecular structure of $(R_C,R_C,R_P/S_C,S_C,S_P)\text{-}[Cp_2Zr(Me)\text{-}(\textit{cyclo}\text{-}1\text{-}O\text{-}2\text{-}PH(Tipp)\text{–}C_6H_{10})]$, 126
Phosphametallacycles
[2 + 2]-Cycloaddukte PH-funktioneller Phosphenium-Komplexe mit Alkylisothiocyanaten: Darstellung von $C_3R_3(OC)_2$<ovl type = "tsqb" > M–P(H)(*t*-Bu)–C(=NR′)–S (R = H, Me; M = Mo, W; R′ = Me, Et, *t*-Bu) und Reaktion unter Beanspruchung der PH-Funktion: Phosphenium-Übergangsmetallkomplexe, 37, 285
Phosphenium complexes
[2 + 2]-Cycloaddukte PH-funktioneller Phosphenium-Komplexe mit Alkylisothiocyanaten: Darstellung von $C_3R_3(OC)_2$<ovl type = "tsqb" > M–P(H)(*t*-Bu)–C(=NR′)–S (R = H, Me; M = Mo, W; R′ = Me, Et, *t*-Bu) und Reaktion unter Beanspruchung der PH-Funktion: Phosphenium-Übergangsmetallkomplexe, 37, 285
Phosphide
Binuclear hydrido platinum(II) complexes: syntheses of $[\{Pt(P\text{–}P)\}_2(\mu\text{-}CHCH_2R)(\mu\text{-}H)][BF_4]$ ($R = C_6H_5$, H) and easy cleavage of a P–C bond in a chelating 1,4-bis (diphenyl)phosphinobutane (P–P), 224
Phosphine
Platinum complexes of heteroannularly bridged heterobidentate ferrocenyl diphosphine ligands: their molecular structure and their use in catalytic carbonylation reactions, 93
Phosphine complexes
Organopalladium(IV) complexes containing phosphine ligands, and the structure of the platinum(IV) complex $[PtMe_3(bpy)(PPh_3)][O_3SCF_3]$ (bpy = 2,2′-bipyridine), 296
Phosphinothiolate complexes
Diorganotin(IV) derivatives of arenephosphinothiol ligands. The crystal structure of $[Ph_2Sn\{2\text{-}(Ph_2P)C_6H_4S\}_2]$ and $[Me_2Sn\text{-}\{2\text{-}(Ph_2PO)\text{-}6\text{-}(Me_3Si)C_6H_3S\}_2]$, 59
Phospholes
Convenient route for the preparation of unsymmetrical phospholes via zirconacyclopentadienes, 261
Phosphorus
The thermal rearrangement of $-\{Me_2Si\text{-}\eta^5\text{-}C_5H_4(CO)Fe(CO)_2\text{-}Fe(CO)\text{-}\eta^5\text{-}C_5H_4SiMe_2\}-$ in the presence of phosphorus ligands, 268

Phosphorus heterocycle
Convenient route for the preparation of unsymmetrical phospholes via zirconacyclopentadienes, 261
Photolysis
Evidence for the involvement of pentacoordinate P-intermediates during the UV light-mediated fragmentation of a 2-phosphabicyclo[2.2.2]octene 2-oxide in the presence of O-nucleophiles, 109
Reactivity of $[Fe_2(CO)_6(\mu\text{-}CO)(\mu\text{-}dppm)]$ towards unsaturated sulfur-containing compounds, SO_2, CS_2 and ArNCS: facile carbon–sulfur bond cleavage at the diiron centre, 134
Platinum
Binuclear hydrido platinum(II) complexes: syntheses of $[\{Pt(P\text{–}P)\}_2(\mu\text{-}CHCH_2R)(\mu\text{-}H)][BF_4]$ ($R = C_6H_5$, H) and easy cleavage of a P–C bond in a chelating 1,4-bis (diphenyl)phosphinobutane (P–P), 224
New cyclometallated platinum(II) compounds with thiosemicarbazones: crystal and molecular structure of $[Pt\{4\text{-}MeC_6H_3C(Me)=NN=C(S)NH_2\}(PPh_3)]$, 199
Organopalladium(IV) complexes containing phosphine ligands, and the structure of the platinum(IV) complex $[PtMe_3(bpy)(PPh_3)][O_3SCF_3]$ (bpy = 2,2′-bipyridine), 296
Platinum complexes of heteroannularly bridged heterobidentate ferrocenyl diphosphine ligands: their molecular structure and their use in catalytic carbonylation reactions, 93
Reactions of $[Pt_2(\mu\text{-}S)_2(PPh_3)_4]$ with Group 6 and 7 metal carbonyls; crystal structure of the apparently unsaturated heterometallic complex $[Pt_2Re(\mu_3\text{-}S)_2(PPh_3)_4(CO)_3]^+$, 276
Platinum complex
The X-ray crystallographic study of the reaction of bis(2-thienyl)ditelluride with tetrakis(triphenylphosphine)-platinum or -palladium, 232
Polycarbodiimido-germylene
Bis(trimesitylgermylcarbodiimido)germylene, trimesitylgermylcyanamide and trimesitylgermylcarbodiimide, 153
Porphyrins
Isocyanide ligation at ruthenium(II) complexes with chelating tertiary amine and porphyrin ligands. Structural and electrochemical studies, 114
Potassium anions
Electron-transfer processes in the reaction of K^-, K^+(15-crown-5)$_2$ with carbazole: unexpected formation of metallic potassium, 66
Pyrazolylborate complexes
Synthesis and structure of Rh(I) tris(pyrazolyl)borate complexes of the type $Tp^{Ph,Me}RhL_1L_2$, 178
Pyridinylpyrazole
Synthesis and structural analysis of palladium(II) pyridinylpyrazole complexes by ^1H-, ^{13}C-, ^{15}N-NMR and X-ray diffraction. Comparison of binuclear methylpalladium, chloromethylpalladium, and dichloropalladium complexes by ^{15}N-NMR, 208

Rearrangement
The thermal rearrangement of –$[Me_2Si\text{-}\eta^5\text{-}C_5H_4(CO)Fe(CO)_2Fe(CO)\text{-}\eta^5\text{-}C_5H_4SiMe_2]$– in the presence of phosphorus ligands, 268
Reversibility
Evidence for the involvement of pentacoordinate P-intermediates during the UV light-mediated fragmentation of a 2-phosphabicyclo[2.2.2]octene 2-oxide in the presence of O-nucleophiles, 109
Rhenium
Reactions of $[Pt_2(\mu\text{-}S)_2(PPh_3)_4]$ with Group 6 and 7 metal carbonyls; crystal structure of the apparently unsaturated heterometallic complex $[Pt_2Re(\mu_3\text{-}S)_2(PPh_3)_4(CO)_3]^+$, 276
Rhodium
Regioselective C–F and C–H bond activation: synthesis and structure of an asymmetric rhodium complex comprising a $\eta^5\text{:}\eta^1\text{:}\eta^1$-cyclopentadienyl-bis(phosphine) ligand, 292
Synthesis and structure of Rh(I) tris(pyrazolyl)borate complexes of the type $Tp^{Ph,Me}RhL_1L_2$, 178
Rhodium-catalyzed phenylation
Rhodium-catalyzed phenylation of *N*-arylsulfonyl aldimines with sodium tetraphenylborate or trimethyl(phenyl)stannane, 31
Ring compounds
cyclo-Boratrisiloxane and *cyclo*-diboratetrasiloxane derivatives and their reactions with amines: crystal and molecular structure of $(p\text{-}BrC_6H_4BO)_2(Ph_2SiO)_2$, 241
Ruthenium
Isocyanide ligation at ruthenium(II) complexes with chelating tertiary amine and porphyrin ligands. Structural and electrochemical studies, 114

Schiff bases
New cyclometallated platinum(II) compounds with thiosemicarbazones: crystal and molecular structure of $[Pt\{4\text{-}MeC_6H_3C(Me)=NN=C(S)NH_2\}(PPh_3)]$, 199
Silanones
Matrix isolation infrared and density functional theoretical studies of cryogenic reactions of silicon atoms with dimethyl ether and methanol: new route to generation and stabilization of transient silanones, 248
Silicon
cyclo-Boratrisiloxane and *cyclo*-diboratetrasiloxane derivatives and their reactions with amines: crystal and molecular structure of $(p\text{-}BrC_6H_4BO)_2(Ph_2SiO)_2$, 241
Regiospecific synthesis of alpha-lithiated alkoxysilanes, 87
The thermal rearrangement of –$[Me_2Si\text{-}\eta^5\text{-}C_5H_4(CO)Fe(CO)_2Fe(CO)\text{-}\eta^5\text{-}C_5H_4SiMe_2]$– in the presence of phosphorus ligands, 268
Silicon atoms
Matrix isolation infrared and density functional theoretical studies of cryogenic reactions of silicon atoms with dimethyl ether and methanol: new route to generation and stabilization of transient silanones, 248
Sodium tetraphenylborate
Rhodium-catalyzed phenylation of *N*-arylsulfonyl aldimines with sodium tetraphenylborate or trimethyl(phenyl)stannane, 31
Stable germylene
Bis(trimesitylgermylcarbodiimido)germylene, trimesitylgermylcyanamide and trimesitylgermylcarbodiimide, 153
Structure determinations
Synthese und Struktur von {{*N*-[Bis(diisopropylamino)phosphanyl]amino(phenyl)carben}dicarbonyl(η^5 - methylcyclopentadienyl)mangan(0)} und versuchte Darstellung von {[2-Bis(trimethylsilyl)methyl-3-phenyl-2*H*-azaphosphiren-κ*P*] dicarbonyl(η^5-methylcyclopentadienyl)mangan(0)}, 12
Structures
Reactions of $[\{\mu\text{-}\eta^2\text{-}C(R^1)=CHR^2\}(\mu\text{-}CO)Fe_2(CO)_6]^-$ with $(\mu\text{-}S_2)Fe_2(CO)_6$: synthesis and structures of $[\mu\text{-}\eta^2\text{-}C(R^1)=CHR^2](\mu\text{-}RS)[Fe_2(CO)_6]_2(\mu_4\text{-}S)$ (R = alkyl, phenyl or acyl), 158
Substitution
The thermal rearrangement of –$[Me_2Si\text{-}\eta^5\text{-}C_5H_4(CO)Fe(CO)_2Fe(CO)\text{-}\eta^5\text{-}C_5H_4SiMe_2]$– in the presence of phosphorus ligands, 268
Sulfide
Reactions of $[Pt_2(\mu\text{-}S)_2(PPh_3)_4]$ with Group 6 and 7 metal carbonyls; crystal structure of the apparently unsaturated heterometallic complex $[Pt_2Re(\mu_3\text{-}S)_2(PPh_3)_4(CO)_3]^+$, 276
Sulfur
Synthesis and spectroscopic characterisation of compounds with formula $[\{Fe_2S_2(CO)_5\}Ph_2P(CH_2)_nPPh_2\{Fe(CO)_4\}]$ and $[\{Fe_2S_2(CO)_5\}_2Ph_2P(CH_2)_nPPh_2]$ (n = 3, 4 and 6), 191

Synthesis, optical nonlinearity and crystal structure of a novel mixed-chalcogen, thiolato-bridged complex $[(\eta^5\text{-}C_5H_5)_2\text{-}Mo_2(\mu\text{-}S)(\mu\text{-}Te)(\mu\text{-}SPH)_2]$, 140

Sulfur dioxide

Reactivity of $[Fe_2(CO)_6(\mu\text{-}CO)(\mu\text{-}dppm)]$ towards unsaturated sulfur-containing compounds, SO_2, CS_2 and ArNCS: facile carbon–sulfur bond cleavage at the diiron centre, 134

Suzuki reaction

N-Heterocyclic carbenes: Part 26. *N*-Heterocyclic carbene complexes of palladium(0): synthesis and application in the Suzuki cross-coupling reaction, 186

Synthesis

Reactions of $[\{\mu\text{-}\eta^2\text{-}C(R^1)\text{=}CHR^2\}(\mu\text{-}CO)Fe_2(CO)_6]^-$ with $(\mu\text{-}S_2)Fe_2(CO)_6$: synthesis and structures of $[\mu\text{-}\eta^2\text{-}C(R^1)\text{=}CHR^2](\mu\text{-}RS)[Fe_2(CO)_6]_2(\mu_4\text{-}S)$ (R = alkyl, phenyl or acyl), 158

Tantalum

Chemical behaviour of alkyl imido cyclopentadienyl niobium and tantalum(V) complexes in insertion processes. X-ray crystal structures of $[MCpCl(NAr)\{\eta^2\text{-}C(Me)\text{=}NAr\}]$ (Ar = 2,6-$Me_2C_6H_3$; M = Nb, Cp = $\eta^5\text{-}C_5H_4SiMe_3$; M = Ta, Cp = $\eta^5\text{-}C_5Me_5$) and $[Ta(\eta^5\text{-}C_5Me_5)Me(NAr)\{\eta^2\text{-}C(CH_2CMe_2Ph)\text{=}O\}]$ (Ar = 2,6-$Me_2C_6H_3$), 36

Tellurium

Synthesis, optical nonlinearity and crystal structure of a novel mixed-chalcogen, thiolato-bridged complex $[(\eta^5\text{-}C_5H_5)_2Mo_2(\mu\text{-}S)(\mu\text{-}Te)(\mu\text{-}SPh)_2]$, 140

Tertiary amine

Isocyanide ligation at ruthenium(II) complexes with chelating tertiary amine and porphyrin ligands. Structural and electrochemical studies, 114

Thienylcopper

Tetrameric thienylcopper and pentanuclear thienylcuprate, 102

Thienylcuprate

Tetrameric thienylcopper and pentanuclear thienylcuprate, 102

Thiolato

Synthesis, optical nonlinearity and crystal structure of a novel mixed-chalcogen, thiolato-bridged complex $[(\eta^5\text{-}C_5H_5)_2\text{-}Mo_2(\mu\text{-}S)(\mu\text{-}Te)(\mu\text{-}SPh)_2]$, 140

Thiosemicarbazone

New cyclometallated platinum(II) compounds with thiosemicarbazones: crystal and molecular structure of $[Pt\{4\text{-}MeC_6H_3C(Me)\text{=}NN\text{=}C(S)NH_2\}(PPh_3)]$, 199

Tin complexes

Diorganotin(IV) derivatives of arenephosphinothiol ligands. The crystal structure of $[Ph_2Sn\{2\text{-}(Ph_2P)C_6H_4S\}_2]$ and $[Me_2Sn\{2\text{-}(Ph_2PO)\text{-}6\text{-}(Me_3Si)C_6H_3S\}_2]$, 59

Trien

Synthesis of dioxazaborocines from *N,N'*-alkylbridged-bis(bis(2-hydroxybenzyl)aminomethyl)amines, 215

Trimesitylgermylcarbodiimide

Bis(trimesitylgermylcarbodiimido)germylene, trimesitylgermylcyanamide and trimesitylgermylcarbodiimide, 153

Trimesitylgermylcyanamide

Bis(trimesitylgermylcarbodiimido)germylene, trimesitylgermylcyanamide and trimesitylgermylcarbodiimide, 153

Trimethyl(phenyl)stannane

Rhodium-catalyzed phenylation of *N*-arylsulfonyl aldimines with sodium tetraphenylborate or trimethyl(phenyl)stannane, 31

Tungsten

Reactions of $[Pt_2(\mu\text{-}S)_2(PPh_3)_4]$ with Group 6 and 7 metal carbonyls; crystal structure of the apparently unsaturated heterometallic complex $[Pt_2Re(\mu_3\text{-}S)_2(PPh_3)_4(CO)_3]^+$, 276

Unsaturated ligand

Convenient routes to mono- and dinuclear (C_5Me_5)iron(II) complexes bearing acyl, alkynyl, aryl and thienyl ligands, 81

X-ray crystallography

The X-ray crystallographic study of the reaction of bis(2-thienyl)ditelluride with tetrakis(triphenylphosphine)platinum or -palladium, 232

X-ray crystal structure determination

Tetrameric thienylcopper and pentanuclear thienylcuprate, 102

X-ray diffraction

Monomeric organoaluminium complexes $RAl(OR^*)_2$ and R_2AlOR^* with an optically active amino alkoxide ligand, 21

Zirconacyclopentadienes

Convenient route for the preparation of unsymmetrical phospholes via zirconacyclopentadienes, 261

Zirconium

Synthesis and characterization of Group 4 metallocene complexes with two disiloxanediyl bridges, 147

Zirconocene complex

P–H functionalised phosphinoalkoxides as ligands: synthesis and catalytic properties of $[Cp_2Zr(Me)(cyclo\text{-}1\text{-}O\text{-}2\text{-}PH(Tipp)\text{-}C_6H_{10})]$ (Tipp = 2,4,6-$Pr^i_3C_6H_2$), $[Cp_2Zr(Me)(cyclo\text{-}1\text{-}O\text{-}2\text{-}PH(Tipp)\{Mo(CO)_5\}\text{-}C_6H_{10})]$ and $[Cp_2Zr(BH_4)(cyclo\text{-}1\text{-}O\text{-}2\text{-}PH(Mes)(BH_3)\text{-}C_6H_{10})]$ (Mes = 2,4,6-$Me_3C_6H_2$), and molecular structure of $(R_C,R_C,R_P/S_C,S_C,S_P)$-$[Cp_2Zr(Me)(cyclo\text{-}1\text{-}O\text{-}2\text{-}PH(Tipp)\text{-}C_6H_{10})]$, 126